	pIMAGO®-biotin Phosphoprotein Detection Kit For Microplate Fluor 550 detection		on Kit
ore	Cat. # 901-400 1 Kit	Orders: Support:	 order@tymora-analytical.com customersupport@tymora-analytical.com 765-490-6834
	(400 wells)	Web:	www.tymora-analytical.com

This product is for *in vitro* research use only and is not intended for use in humans or animals.

Introduction.

pIMAGO is a universal phosphoprotein detection technology that enables sensitive and specific recognition of phosphorylated molecules. Unlike phospho-antibodies, the binding is not biased by amino acid sequence, and therefore can be used for detection of any phosphorylation event on any protein site. pIMAGO detection protocol resembles a simple ELISA procedure and can be easily incorporated by any laboratory. Due to its small size, pIMAGO can be multiplexed with antibodies for simultaneous detection of phosphorylation and total protein amount.

Results.

pIMAGO-based detection and quantitation of phosphorylated b-casein signal compared to nonphosphoprotein BSA (left figure). Multiplexed detection of phospho-Cdb3 using pIMAGO in the 700 channel and anti-Cdb3 antibody in the 800 channel (right figure). pIMAGO-based kinase assays of 5 kinases and their substrates, including control, ATP, and ATP + Kinase wells (bottom figure).


pIMAGO colorimetric detection



Phosphorylated b-casein (1-100ng)



pIMAGO fluorescence-based detection



pIMAGO detection of in vitro kinase assays

■ Control ■ ATP ■ ATP + Kinase 2.5 Sample OD415 2 1.5 1 0.5 0 Abl Cdk1 Plk1 Syk BRAF

Original citation.

3

Iliuk A, Martinez J, Hall MC, Tao WA (2011). Phosphorylation assay based on functionalized soluble nanopolymer. Anal. Chem. 83(7): 2767-74.

PMID: 21395237

Kit components.

- 400µL of pIMAGO reagent \checkmark
- 400µL of avidin-Fluor550
- 40mL of Binding buffer
- 240mL of Blocking buffer
- 240mL of pIMAGO buffer
- 200µL of control phosphoprotein
- 10 flat-bottom clear 96-well plates